

KOPI

D11

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 31/65, 47/02, 47/10, 47/22, 47/00	A1	(11) International Publication Number: WO 96/01634 (43) International Publication Date: 25 January 1996 (25.01.96)
(21) International Application Number: PCT/GB95/01583 (22) International Filing Date: 5 July 1995 (05.07.95) (30) Priority Data: 9413873.2 9 July 1994 (09.07.94) GB (71) Applicant (for all designated States except US): NORBROOK LABORATORIES LIMITED [GB/GB]; Station Works, Camlough Road, Newry BT35 6JP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): PATTERSON, Alan [GB/GB]; 60 Quarry Road, Belfast BT4 2NQ (GB). HOLMES, Drew [GB/GB]; 9 Village Manor, Bryansford, Newcastle, Co Down BT33 0SL (GB). (74) Agent: FITZPATRICKS; 4 West Regent Street, Glasgow G2 IRS (GB).		(81) Designated States: AU, CA, CN, GB, JP, MX, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: LONG-ACTING OXYTETRACYCLINE COMPOSITION		
(57) Abstract An injectable composition of higher residual effect with reduced detrimental effects such as pain at injection site, swelling, tissue irritancy or necrosis and containing as active principle a tetracycline compound, either as the free base or a salt thereof with a physiologically acceptable acid, complexed with a substantially equimolar amount of a magnesium compound, is solubilised in a water miscible solvent system comprising, either (i) a) glycerol formal in an amount of from about 10 to about 50 % v/v; with b) polyethylene glycol in an amount of from about 1 to 15 % v/v; or (ii) from about 25 to about 75 % v/v of N-methylpyrrolidone, said composition optionally containing a pH modifier in an amount sufficient to maintain a physicochemically acceptable pH, the balance being made up with water q.s.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Long-Acting Oxytetracycline Composition

This invention relates to injectable formulations containing tetracyclines, particularly oxytetracycline, which exhibit higher residual effect with less of the known detrimental effects such as pain at injection site, swelling, tissue irritancy or necrosis.

Preparation of pharmaceutical compositions containing tetracyclines and oxytetracycline in particular has always presented a challenge due to aqueous solubility constraints which firstly have impact upon composition stability, and secondly upon parenteral administration.

Prior art oxytetracycline compositions have exhibited relatively high viscosity at low temperatures which makes injection difficult, have shown poor stability and suffered limitations on strength of active principle. Thus considerable research has gone into determining suitable complexing agents and more favourable co-solvents to address these shortcomings. A review of the art suggests that presence of calcium, and especially magnesium in the formulation now appears mandatory as a complexing agent and whereas some improvements have been made in stability and delivery by adopting various co-solvent systems, higher concentration loadings and residual effect remain areas in which improvements are needed. This is especially of interest for veterinary purposes where the need is to deliver high effective doses with minimum effort in animal handling and detrimental effect on the animal requiring treatment.

At the current time prior art so-called "long-acting" oxytetracycline formulations typically contain 200 mg/ml oxytetracycline and are administered at 20 mg/kg body weight, having activity as determined by residual blood levels of oxytetracycline detectable for up to about four days or so.

An object of this invention is to provide a composition of substantially greater long acting effect whilst minimising to the greatest extent possible the defects observed in previously proposed formulations. In particular the invention to be particularly described hereinbelow

provides for administration of an oxtetracycline formulation at dose rates of from 10 to 40 mg/kilogram bodyweight, giving at 30 mg/kg in animals an extended duration of effective plasma levels against susceptible organisms in excess of 9 days which is a surprising achievement in the light of the known prior art.

Solubility of oxytetracycline in non-aqueous solvents was considered by Eugene Gans and Takeru Higuchi, Journal of the American Pharmaceutical Association, 1957, Vol XLVI, pp 587-591.

The patent literature in this area is extensive and one could refer to the following patents which are illustrative of the decades of research carried out on formulation of tetracycline compositions:

GB-A-894 619, GB-A-1 131 007, GB-A-1 250 304, GB-A-1 286 351, GB-A-1 427 882, GB-A-1 494 558, GB-A-1 508 601, GB-A-1 514 838, GB-A-1 520 197, GB-A-1 538 903 GB-A-1 563 478, GB-A-1 592 053, GB-B-2 047 097; EP-B-38 103, EP-B-96 942; US-A-2 516 080, US-A-2 980 584, US-A-2 990 331, US-A-3 062 717, US-A-3 219 529, US-A-3 557 280, US-A-3 712 949, US-A-3 957 972, US-A-4 011 313, US-A-4 018 889, US-A-4 020 162, US-A-4 126 680, US-A-4 386 083, US-A-4 399 127, US-A-4 772 460, US-A-4 957 972, and US-A-5 075 295.

From these documents it is apparent that a variety of water-dispersible complex-stabilisers or water-miscible co-solvents have been proposed including 2-pyrrolidone, polyvinyl pyrrolidone, polyethylene glycols, caprolactam, 2-piperidone, and glycerol formal (a reaction product of glycerol and formaldehyde) in specific formulations. However it is by no means clear that the said co-solvents are equally interchangeable nor can the effect of such a change be entirely predictable for a given formulation.

US-A-4 386 083 proposes use of glycerol formal in conjunction with magnesium acetate and magnesium chloride, whilst US-A-4 772 460 proposes use of N-methylpyrrolidone (1-methyl-2-pyrrolidone) and a soluble magnesium compound.

US-A-5 075 295 is particularly directed to a composition aiming to achieve up to 30% oxytetracycline, which contains polyethylene glycol 400 and magnesium oxide, but examples given only appear to show a capability of achieving up to 5 25% oxytetracycline and there is to applicant's knowledge no current commercially available product capable of achieving greater than 20%.

Accordingly this invention provides a composition containing as active principle a tetracycline compound, 10 either as the free base or a salt thereof with a physiologically acceptable acid, complexed with a substantially equimolar amount of a magnesium compound, solubilised in a water miscible solvent system comprising, either

15 (i) a) glycerol formal in an amount of from about 10 to about 50% v/v; with
b) polyethylene glycol in an amount of from about 1 to 15% v/v; or

(ii) from about 25 to about 75% v/v of N-methylpyrrolidone, 20 said composition optionally containing a pH modifier in an amount sufficient to maintain a physiochemically acceptable pH, the balance being made up with water q.s.

The composition optionally contains a thickener such as polyvinyl pyrrolidone in an amount of up to 10% w/v, and may 25 contain usual formulation aids or auxiliaries typically used for such formulations. Thus the composition may contain antioxidants, e.g. sodium formaldehyde sulfoxylate and pH adjusting agents e.g. monoethanolamine, to provide a preferred pH range of from about 7.5 to about 9.5, more 30 preferably from about 8.5 to about 9.0.

Preferably the composition contains a magnesium compound such as magnesium oxide or a salt e.g magnesium chloride.

The preferred compositions contain oxytetracycline as 35 the base or its hydrochloride in an amount of from about 15 to about 35% w/v, complexed with an equimolar ratio of a magnesium compound, preferably a salt, solubilised in a solvent system comprising polyethylene glycol in an amount

of from about 1 to about 15% v/v and glycerol formal in an amount of from about 10 to about 50% v/v. In particular the most preferred composition contains about 30% w/v oxytetracycline, about 40% glycerol formal, about 10% v/v polyethylene glycol with a magnesium-containing complexing agent or stabiliser, antioxidant and water making up the balance. In that composition magnesium oxide is suitably present in an amount of about 2.7% w/v and, as antioxidant, sodium formaldehyde sulphonylate in an amount of about 0.4% w/v may be used. Thus according to the present invention there is provided a formulation capable of providing from about 10 to about 40 mg/kg bodyweight consisting of:

	Oxytetracycline	300 mg
	Magnesium oxide	27 mg
15	Sodium formaldehyde sulphonylate	4 mg
	Glycerol formal	0.4 ml
	Polyethylene glycol	0.1 ml
	Monoethanolamine	q.s. pH 8.6 to 8.8
	Water for injections	<u>to</u> 1 ml

20 The invention will now be further described by way of example for the purposes of practical illustration only.

An oxytetracycline formulation was prepared according to the procedure indicated below using the following components:

25 Active Ingredient -

Oxytetracycline	30% w/v
-----------------	---------

Excipients -

	Magnesium oxide	2.7% w/v
	Sodium formaldehyde sulphonylate	0.4% w/v
30	Glycerol formal	40% w/v
	Polyethylene glycol	10% w/v
	Monoethanolamine	q.s. pH 8.6 to 8.8
	Water for injections	<u>to</u> 100% w/v

35 A controlled environment having an inert atmosphere was provided within which suitable mixing and temperature controllable heating apparatus was assembled. A nitrogen blanket is considered suitable for this purpose. The above

components of the proposed composition were mixed by initially mixing a proportion of the total water with the selected solvents. The sodium formaldehyde sulphoxylate, magnesium oxide and oxytetra-cycline were added sequentially whilst mixing continuously and maintaining a temperature of approximately 65°C until all the constituents have dissolved. Thereafter, the composition is cooled to below 30°C and the pH is adjusted to lie within the range of 8.0 to 9.0, in this case by adding a sufficient amount of mono-ethanolamine. Finally the volume is made up with water, the pH checked and adjusted if necessary, and the composition is filtered through a 0.2 µm filter and filled into appropriate containers.

In alternative embodiments, where use of a thickener such as polyvinyl pyrrolidone is called for then it should preferably be added after the sodium formaldehyde sulphoxylate.

The following Tables provide details of Examples 1 to 14 each of which provided compositions showing excellent stability and which achieved the desired dosage levels and long acting effect.

TABLE I:

INGREDIENTS	EXAMPLE									
	1	2	3	4	5	6	7	8	9	10
Oxytetracycline (% w/v)	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	15.0	35.0
Magnesium Oxide (% w/v)	2.7	2.7	2.7	2.7	2.7	2.7	2.7	13.25*	1.3	3.06
Sodium Formaldehyde Sulphoxylate (% w/v)	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.4	0.4
Glycerol Formal (% v/v)	30.0	30.0	30.0	35.0	35.0	35.0	40.0	40.0	40.0	40.0
Polyethylene Glycol 200 (% v/v)	10.0	15.0	20.0	10.0	15.0	20.0	10.0	10	10.0	10.0
Polyvinyl Pyrrolidone K12 (% w/v)			3.0							
Water to (% v/v)	100	100	100	100	100	100	100	100	100	100

*Magnesium Chloride

TABLE 2:

INGREDIENTS	EXAMPLE			
	11	12	13	14
Oxytetracycline (% w/v)	30	30	25	35
Magnesium Oxide (% w/v)	2.78	2.78	2.3	3.21
N-Methyl Pyrrolidone (% v/v)	30.0	60.0	60.0	60.0
Sodium Formaldehyde Sulphoxylate (% w/v)	0.4	0.4	0.40	0.4
Water to (% v/v)	100	100	100	100

Claims

1. A composition containing as active principle a tetracycline compound, either as the free base or a salt thereof with a physiologically acceptable acid, complexed with a substantially equimolar amount of a magnesium compound, solubilised in a water miscible solvent system comprising, either
- (i) a) glycerol formal in an amount of from about 10 to about 50% v/v; with
- b) polyethylene glycol in an amount of from about 1 to 15% v/v; or
- (ii) from about 25 to about 75% v/v of N-methylpyrrolidone, said composition optionally containing a pH modifier in an amount sufficient to maintain a physiochemically acceptable pH, the balance being made up with water q.s.
2. A composition according to claim 1 comprising as a thickener polyvinyl pyrrolidone in an amount of up to about 10% w/v.
3. A composition according to claim 1 or claim 2 wherein the magnesium compound is magnesium oxide.
4. A composition according to claim 1 or claim 2 wherein the magnesium compound is a magnesium salt.
5. A composition according to claim 4 wherein the magnesium salt is magnesium chloride.
6. A composition according to claim 1 wherein the tetracycline compound is oxytetracycline base or its hydrochloride in an amount of from about 15 to about 35% w/v.

7. A composition according to claim 1 wherein the composition contains about 30% w/v oxytetracycline, about 40% glycerol formal, about 10% v/v poly-ethylene glycol with a magnesium-containing complexing agent or stabiliser, antioxidant and water making up the balance.

8. A composition according to claim 7 wherein magnesium oxide is present in an amount of about 2.7% w/v and, as antioxidant, sodium formaldehyde sulfoxylate in an amount of about 0.4% w/v may be used.

9. An injectable composition for treatment of animals which consists of:

15	Oxytetracycline	300 mg
	Magnesium oxide	27 mg
	Sodium formaldehyde sulfoxylate	4 mg
	Glycerol formal	0.4 ml
	Polyethylene glycol	0.1 ml
	Monoethanolamine	q.s. pH 8.6 to 8.8
20	Water for injections	to 1 ml, the

said composition providing for administration of from about 10 to about 40 mg of oxytetracycline per kilogram of bodyweight.

25 10. An injectable composition for treatment of animals according to any one of the Examples 1 to 14 hereinbefore.

INTERNATIONAL SEARCH REPORT

Int ional Application No

PCT/GB 95/01583

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 A61K31/65 A61K47/02 A61K47/10 A61K47/22 A61K47/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 096 942 (AESCULAAP B.V.,NL) 28 December 1983 see claims see examples I-V ---	1-3
X	US,A,4 772 460 (S.U.MALOOK ET AL.) 20 September 1988 cited in the application see claims see examples ---	1-3,5
Y	EP,A,0 038 013 (DIAMOND SHAMROCK CORP.,U.S.A.) 21 October 1981 cited in the application see the whole document --- -/--	1-10
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 7 November 1995		Date of mailing of the international search report 17.11.95
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016		Authorized officer Scarponi, U

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/GB 95/01583

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DE,A,20 11 793 (DIAMOND SHAMROCK CORP.,U.S.A.) 17 September 1970 see the whole document ----	1-10
Y	US,A,5 075 295 (J.A.ZUPAN ET AL.) 24 December 1991 cited in the application see the whole document -----	1-10
Y	FR,A,2 081 434 (PFIZER CORP.) 3 December 1971 see the whole document -----	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/01583

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0096942	28-12-83	NL-A- 8202423	02-01-84
US-A-4772460	20-09-88	BE-A- 905528	02-02-87
EP-A-0038013	21-10-81	CA-A- 1163197	06-03-84
		US-A- 4399127	16-08-83
		US-A- 4386083	31-05-83
DE-A-2011793	17-09-70	BE-A- 747255	14-09-70
		FR-A, B 2034890	18-12-70
		GB-A- 1250304	20-10-71
		NL-A- 7003555	15-09-70
		OA-A- 3376	15-12-70
		US-A- 3712949	23-01-73
US-A-5075295	24-12-91	NONE	
FR-A-2081434	03-12-71	DE-A- 2001604	19-08-71
		GB-A- 1286351	23-08-72